Hip Fracture Deaths in Utah Hospitals (IQI 19)

Adults 18 Years and Over: 2004-2006

	Total Hip		T . 1	C 1
TT	Fracture	Actual	Expected	
Hospital	Patients	Deaths	Deaths	Rating
Utah Overall	4,973	3.0%	3.0%	**
LDS Hospital	407	0.7%	3.0%	***
Alta View Hospital	197	3.0%	4.2%	**
American Fork Hospital	170	3.5%	2.5%	**
Ashley Regional Medical Center	45	6.7%	2.5%	**
Brigham City Community Hospital	74	2.7%	2.2%	**
Castleview Hospital	81	1.2%	3.4%	**
Cottonwood Hospital	270	2.2%	4.3%	**
Davis Hospital	65	1.5%	1.5%	**
Dixie Regional Medical Center	460	2.4%	2.5%	**
Lakeview Hospital	211	2.4%	2.8%	**
Logan Regional	192	4.2%	3.7%	**
McKay-Dee Hospital	402	3.5%	3.4%	**
Mountain View Hospital	109	4.6%	3.3%	**
Mountain West Hospital	62	3.2%	4.5%	**
Ogden Regional Medical Center	172	2.3%	2.3%	**

Most Utah hospitals in this table had about as many deaths as expected compared to similar hospitals in the U.S. One Utah hospitals had fewer than expected. Two hospitals had more than expected. Hospitals that had no deaths of these patients during this time period also received a three-star rating.

The following hospitals treated at least one patient but less than 30 patients:

Allen Memorial Hospital, Bear River Valley Hospital, Beaver Valley Hospital, Cache Valley Specialty Hospital, Central Valley Hospital, Gunnison Valley Hospital, Heber Valley Hospital, Primary Children's Medical Center, Sanpete Valley Hospital, Sevier Valley Hospital and The Orthopedic Specialty Hospital.

Continued on Next Page

Statistical Rating: based on the Exact 95% Confidence Interval test of statistical significance, except hospitals with no deaths from 2004 through 2006 have three stars. Rating based on small numbers may be unreliable.

NOTE: Hospitals are listed alphabetically within each star rating.

Click here to read Data Limitations.

Actual deaths: percentage of patients who received care for this condition or procedure and died in this hospital.

Expected deaths: percentage of patients who were expected to die at this hospital if it performed the same as other U.S. hospitals that treated similar patients (age, gender, how ill the patients were).

Hip Fracture Deaths in Utah Hospitals (IQI 19)

Adults 18 Years and Over: 2004-2006

- CONTINUED -

	Total Hip Fracture		Expected	Statistical
Hospital	Patients	Deaths	Deaths	Rating
Utah Overall	4,973	3.0%	3.0%	**
Orthopedic Specialty Hospital	31	3.2%	1.6%	**
Pioneer Valley Hospital	74	4.1%	2.3%	**
Salt Lake Regional	39	2.6%	2.2%	**
St. Mark's Hospital	581	2.2%	2.7%	**
Uintah Basin Hospital	48	2.1%	1.9%	**
University Health Care	330	2.4%	2.1%	**
Utah Valley Regional Medical Center	472	4.0%	3.1%	**
Valley View Hospital	150	2.0%	1.4%	**
Veterans Administration Medical Center	101	5.9%	5.2%	**
Jordan Valley Hospital	63	7.9%	2.2%	*
Timpanogos Regional Hospital	94	10.6%	4.3%	*

Most Utah hospitals in this table had about as many deaths as expected compared to similar hospitals in the U.S. One Utah hospitals had fewer than expected. Two hospitals had more than expected. Hospitals that had no deaths of these patients during this time period also received a three-star rating.

The following hospitals treated at least one patient but less than 30 patients:

Allen Memorial Hospital, Bear River Valley Hospital, Beaver Valley Hospital, Cache Valley Specialty Hospital, Central Valley Hospital, Gunnison Valley Hospital, Heber Valley Hospital, Primary Children's Medical Center, Sanpete Valley Hospital, Sevier Valley Hospital and The Orthopedic Specialty Hospital.

Statistical Rating: based on the Exact 95% Confidence Interval test of statistical significance, except hospitals with no deaths from 2004 through 2006 have three stars. Ratings based on small numbers may be unreliable.

NOTE: Hospitals are listed alphabetically within each star rating.

Click here to read Data Limitations.

Actual deaths: percentage of patients who received care for this condition or procedure and died in this hospital.

Expected deaths: percentage of patients who were expected to die at this hospital if it performed the same as other U.S. hospitals that treated similar patients (age, gender, how ill the patients were).